



LEAP Report

MRT[®] Food Sensitivity Test Results

And

LEAP ImmunoCalm[®] Dietary Management Program

Prepared For: **SAMPLE PATIENT**
Physician: **WELLNESS, MARK MD**
Date: **12/06/2017**
Identifier: **S70120**










Laboratory Director / Supervisor

Physician: WELLNESS, MARK MD
 Patient: SAMPLE PATIENT
 Identifier: S70120
 Profile: MRT 170
 Test Date: 12/06/2017
 Technologist: JW



Oxford Clinical Laboratory
 3555 Fiscal Court, Suite #9
 Riviera Beach, FL 33404
 Phone: (561) 848-7111
 FL License #: 800027063
 CLIA ID #: 10D0914874
 U.S. Patents: 6,114,174 6,200,815





MRT Results for S. PATIENT, page 2 of 2

Physician: **WELLNESS, MARK MD**
 Patient: **SAMPLE PATIENT**
 Identifier: **S70120**
 Profile: **MRT 170**
 Test Date: **12/06/2017**
 Technologist: **JW**



Oxford Clinical Laboratory
 3555 Fiscal Court, Suite #9
 Riviera Beach, FL 33404
 Phone: (561) 848-7111
 FL License #: 800027063
 CLIA ID #: 10D0914874
 U.S. Patents: 6,114,174 6,200,815



Degrees of reactivity may not in all cases correlate with levels of clinical reactivity to the tested food or chemical. It is recommended to completely avoid all tested items with "Reactive," or "Moderately Reactive" scores. If tested "Non-Reactive" foods have been consumed regularly before testing, there is a high probability that they are "safe" and are not likely to provoke symptoms, as long as there is no history of allergic, autoimmune, or other type of inflammation-provoking or symptom-provoking reaction. These test results can form the basis of an eating plan developed under the guidance of a physician, dietitian, or other qualified healthcare practitioner.